



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**FURTHER NOTES ON THE EMBRYONIC WHORLS OF THE  
MURICIDÆ.**

BY FRANK C. BAKER.

In 1890 I described in these Proceedings (pp. 66-72) the embryonic whorls of a number of species of this most interesting family; and in the Proceedings of the Rochester Academy of Science (Vol. 1, 1891, pp. 129-133) I added several species to those already described. From that time until a few months ago, I was unable to find perfect material in any collection under my charge. Some months since, however, I discovered perfect specimens of *Murex troscheli* and *Eupleura caudata*, with the nuclei intact. Descriptions of these are given below:—

**Murex troscheli**, Lischke. (Fig. 1.)

The nucleus consists of two and a half smooth, glossy, hyaline whorls, gradually increasing in size; a carina begins at the apex and encircles the base of the whorls near the suture below; it terminates




Fig. 1.

in the fourth spiral line of the after growth. There is a considerable varix at the junction of the nucleus with the after growth. The whorls succeeding the nucleus are crossed by four spiral, beaded lines; spinose varices begin on the fourth whorl.

The only species whose nucleus approaches that of the present species is *Murex pliciferus* (Sowb.), but that is higher in proportion to its width and somewhat differently shaped. The number of whorls are the same in both species.

It may not be out of place to indicate here the species of which the nuclei have been described and figured. Descriptions of these may be found in the Proceedings spoken of above. The species described are as follows:—

*Murex scolopax*, Dillw.  
“ *occa*, Sowb.  
“ *tribulus*, Linn.  
“ *rectirostris*, Sowb.

*Murex recurvirostris*, Sowb.  
“ *similis*, Sowb.  
“ *tryoni*, Hidalgo.  
“ *cailleti*, Petit.

<i>Murex brevispina</i> , Lam.	<i>Murex aduncospinosus</i> , Beck.
“ <i>nigrispinosus</i> , Reeve.	“ <i>brandaris</i> , Linn.
“ <i>ternispina</i> , Lam.	“ <i>rufus</i> , Lam.
“ <i>tenuispina</i> , Lam.	“ <i>pliciferus</i> , Sowb.
“ <i>tribulus</i> , Linn.	“ <i>brevifrons</i> , Lam.

**Eupleura caudata**, Say. (Fig. 2.)

The nucleus of this species consists of one and a half semitransparent, glossy whorls; there is no indication of a carina upon them; the tip of the first whorl is immersed in the succeeding whorl and is bent down to one side; the first half of the whorl, looking at the lateral outline, is about half the size of the portion succeeding it; the whorls are well-rounded and full; the sculpture, consisting of spiral and longitudinal lines, begins very faintly on the latter part of the second whorl and gradually grows stronger as the shell increases in size; there



Fig. 2.

are about fourteen longitudinal lines to each whorl, after leaving the nucleus, which are crossed by two heavy spiral lines.

I have seen four specimens of this species, and the characters expressed above appear to be tolerably constant. The present species is the only one of this genus, which I have been able to examine. It will be interesting to know just how variable or constant the nuclei are in the different species. This is a field which needs a great amount of original investigation.